

Lost in music

Lauren Stewart on amusia, and the window it provides on musical development

Listening to music is effortless and highly rewarding for most of us. But people with congenital amusia have life-long difficulties in this regard – they don't recognise tunes that would be familiar to the rest of us and they can't tell one song from another. This article explores why music is so impenetrable to these people and explains how such individuals offer a unique window on to the typical development of the musical brain.

questions
How might congenital amusia affect the understanding of speech in tonal language cultures?

resources
Stewart, L. (2008). Fractionating the musical mind: Insights from congenital amusia. *Current Opinion in Neurobiology*, 18, 127–130.
www.delos.com/listening/home.html
www.goldsmiths.ac.uk/music-mind-brain

references
Cuddy, L.L., Balkwill, L.L., Peretz, I. & Holden, R.R. (2005). Musical difficulties are rare: A study of 'tone deafness' among university students. *Annals of the New York Academy of Sciences*, 1060, 311–324.
Foxton, J.M., Dean, J.L., Gee, R. et al. (2004). Characterization of deficits in pitch perception underlying 'tone deafness'. *Brain*, 127, 801–810.
Hyde, K., Lerch, J.P., Zatorre, R. et al.

Learning that cognitive faculties can go awry, or even fail to develop, gives us special insight into abilities we often take for granted. The capacity to make sense of musical sound is one such example. In Western culture, someone would not count themselves as 'musical' unless they could boast prowess on a musical instrument. But the real prowess, without which musical performance could not develop, is the ability to make sense of musical sound. Music is not 'out there' in the world but must be constructed by our brains. Neural assemblies process pitch, timing and texture, keep track of interwoven melodic lines, and interpret incoming information in the context of the musical rules of our culture.

For most of us, this feat of sophisticated processing is not only effortless but highly rewarding. In contrast to this, a small minority of individuals are denied access to a musical world. Those with a disorder termed congenital amusia have normal hearing and are intellectually unimpaired but lack the implicit knowledge that allows the rest of us to spot 'wrong' notes, recognise familiar tunes from our culture or to tell one tune from another. Such individuals may find music impenetrable and confusing and go to great lengths to avoid it.

Interestingly, there is little overlap between these individuals and those who view themselves as 'tone-deaf' (Cuddy et al., 2005; Wise & Sloboda, 2008). While self-professed 'tone-deaf' individuals have problems that are restricted to musical production (not perception), those with

amusia have a primary perceptual deficit that extends to singing because they cannot use auditory feedback as a guide for musical production.

Those with congenital amusia provide a window on musical development because, by characterising the ways in which musical perception can be compromised, we can start to build a picture of how the normal brain makes sense of musical sound. The study of disordered musical development forms part of a broader research endeavour that draws on experimental psychology, neuropsychology and neuroimaging in order to understand the cognitive neuroscience of music.

A hitch with pitch

Until recently, a problem in making sense of music was regarded as nothing more than an anecdotal curiosity. Some notable figures remarked upon their musical difficulties, including economist Milton Friedman, the revolutionary Che Guevara and former US president Ulysses S. Grant ('I know of only two tunes: one of them is Yankee Doodle Dandy and the other isn't').

The Montreal Battery for the Evaluation of Amusia (MBEA), developed by Isabelle Peretz at the University of Montreal, provides a way to empirically assess musical perception. Listeners discriminate pairs of musical phrases that are either identical or slightly different. In one subtest, the difference involves a note from the original phrase being replaced with a note from a different key; another involves a rearrangement of the timing of the original phrase. Individuals are classified as amusic when their scores fall outside the normal limits of performance – ascertained by testing several hundred individuals to determine a distribution of scores (Peretz et al., 2003).

Identifying individuals who are amusic according to the MBEA is just the first step. A major goal of the research is to understand what the core underlying deficit(s) could be in this musical disorder.

So far, the problem appears to be one of pitch processing: those with amusia have particular difficulty in determining the direction of a pitch change. As a group, those with congenital amusia require pitch changes to exceed one semitone before the direction of pitch change can be successfully resolved (Foxton et al., 2004). One semitone is a common step size in Western music, meaning that those with congenital amusia are able to register pitch changes but are effectively deaf to the pattern of ups and downs ('contour') that makes a melody distinctive.

A problem that may result from this pitch discrimination problem, or exist in parallel with it, involves the capacity to hold on to music in memory. One amusic individual we have worked with complained of her inability to hold music in her head. This is a particular problem given that she is required to sing plainchant five times a day:

When the music finishes the sound is gone – as though it has never happened. And this bewilders me with a sense of failure, to hold on to what I just heard. Others tell me if I try to remember I will, but I never do. I have no idea what people mean when they say, 'I have a tune going round in my head'. I have never had a tune tell out its music in my head, let alone repeat itself!

The perception of music depends upon the integration of events as they unfold in time. It seems that, for this amusic woman, music does not unfold in a continuous stream but is heard as a series of disconnected auditory 'snapshots'. It is interesting that she contrasts her lack of music with the often-reported phenomenon that music 'sticks' in the head. While the 'tune in the head' phenomenon has so far defied explanation, it is tempting to suggest that a lack of music in the head and exuberant musical imagery may reflect opposite ends of the same continuum.

When the feeling's gone

Nabokov, writing in his autobiography, *Speak, Memory* (1966), complains: 'Music, I regret to say, affects me merely as an arbitrary succession of more or less irritating sounds. Under certain emotional circumstances I can stand the spasms of a rich violin, but the concert piano and all wind instruments bore me in small doses and flay me in larger ones. (pp.35–36)



Family case study

A family-run music shop seems an unlikely place to study disordered musical perception. But John O'Neill's shop in Gortin, County Tyrone is where much of our recent research has taken place. His daughter, Anne, read an article I had written about congenital amusia and persuaded three generations of her family to take an online version of the Montreal Battery for the Evaluation of Amusia. Remarkably, half of them scored in the amusic range.

Even though Anne's father, John, struggles to make music himself, he always had musical aspirations for his children. More than 40 years ago, he bought his first son, Sean, a single-row melodian for Christmas and arranged for a local musician to show him the ropes. Before long, friends and neighbours caught on to the idea and sent their own children along to learn. But the interest exceeded the availability of instruments and, to keep the aspiring musicians supplied with fiddles, flutes and accordions, John started sourcing them from all over the world, resulting in the family business he runs today.

While music has become an enormous part of Sean's life, Anne, and several of her siblings, have struggled, despite years of accordion and Irish dancing lessons. 'I desperately wanted to be in a church choir and to be good at singing', she says. 'But every time I turned up the choirmaster said – "Okay, Anne, we have enough children now". I was terribly upset. I knew that however hard I tried, I would never be able to sing like the others. The teacher used to say, "Anne, why don't you play the tambourine?". But because Anne's problem also extended to rhythm, she couldn't keep in time.'

Anne's musical difficulties were readily apparent to her, owing to the musically enriched environment in which she grew up. But because the emphasis was always one of involvement rather than achievement, Anne was always able to derive pleasure from being around music. 'Music brings great joy to us – it's an expression of our culture,' she says. 'My parents celebrated their golden wedding anniversary recently with a special mass. The grandchildren brought a flute to the altar as a symbol of how much music means to our family.'

though these subgroups are apparently similar in their perceptual abilities, they could differ on a perceptual dimension that is not captured by the MBEA. The processing of timbre – the quality of a musical sound – may be compromised in some, but not other, amusics, which may explain the variability in the musical experience. Some indications for timbral, as well as pitch-based deficits come from the comments of several amusics, who report that music sounds like 'banging' or 'noise'.

Family affair

There is good evidence that amusia is a developmental disorder that is under genetic control. A recent study showed that 39 per cent of first degree relatives of those with amusia were similarly affected, compared with only 3 per cent of relatives

of a non-amusic control group (Peretz et al., 2007).

Of course, it can be argued that amusic parents may raise their children in the absence of music, which would also result in inheritance of a musical perceptual disorder. But there are several arguments against such a view. First, infants have been shown to display a high level of musical sensitivity that amusic adults lack, suggesting that the amount of exposure required to develop musically is minimal. Second, if an absence of musical input were a factor, this cannot explain why some siblings within a generation may develop normally. The family case study described in the box on the previous page shows that congenital amusia can be present even in the context of a musically enriched environment, with many years of formal and informal musical training. Importantly, in this family, individuals with amusia coexist with siblings who display heightened levels of musical ability and engagement.

Investigation into pedigrees of amusia set the stage for elucidating candidate genes that may underlie amusia. Brain

imaging studies (Hyde et al., 2007; Mandell et al., 2007) have found subtle differences in frontal and temporal cortical structure, and researchers have hypothesised that these findings reflect an under-connectivity between these two brain areas. Any candidate genes may be responsible for neuronal migration between these two areas in early development.

Conclusion

The study of disordered musical development sets in sharp relief the perceptual abilities that most of us take for granted. By investigating why musical perception fails to develop, we can start to build up a model of the components of normal musical perception. But the study of amusia also has broader significance: researchers are, for instance, using the disorder to ask whether pitch processing in music and language rely on shared or distinct neural resources and whether the pattern processing involved in musical perception bears any relation to other kinds of higher-order relational

processing, both within, and outside the auditory domain.

For those with amusia, the recent focus of attention on the disorder is welcome: the likely biological origins of the disorder provide an account of the difficulties they were previously unable to explain. As researchers learn more about what amusic individuals can and can't perceive, there is always the possibility that this knowledge will help them gain access to a musical world. Just as deaf people learn to enjoy music by listening to its vibrations, those with amusia may be able to gain insight into their residual listening abilities. Many amusics can hear rhythm, and some can also dance: if the 'banging' of Rachmaninov doesn't make it onto their iPod, perhaps drum and bass could set their feet tapping.



I Lauren Stewart
is a Senior Lecturer and Director of a new MSc Music Mind and Brain in Psychology at Goldsmiths, University of London. l.stewart@gold.ac.uk

MA or Diploma in Human Givens Psychotherapy



"A truly inspirational learning experience! I feel privileged to have attended. The tutors' level of knowledge and understanding is remarkable. The human givens approach is creative, eclectic, needs-led and applicable to a wide range of situations."

EDUCATIONAL PSYCHOLOGIST

**MINDFIELDS
COLLEGE**
PSYCHOLOGY • PSYCHOTHERAPY • EDUCATION

Gain proven skills which will make you more effective when treating depression, anxiety-disorders (including PTSD, OCD and phobias), addictions, anger, eating disorders, psychosis, relationship problems, chronic pain and self-harming

THE human givens approach is based on a solid basis of fundamental research and new scientific insights into innate human needs, mind/body functioning, behaviour and psychology. It combines this new knowledge with the most effective, evidence-based psychotherapeutic techniques drawn from a wide range of approaches (including CBT). This practical course rigorously incorporates theory, demonstration and practice. Graduates leave with an in-depth knowledge of the causes of emotional distress and behavioural problems, and the ability to tailor effective psychological interventions to each individual's bio-psycho-social needs.

- Students are taught a powerful package of brief therapy interventions that outcome-informed practice shows are consistently effective with a wide range of conditions
- The tutors counsel patients in front of students to demonstrate the skills of lifting depression/trauma, anxiety management etc.
- The course integrates recent findings about brain functioning which unify person-centred, CBT and solution-focused approaches, lifting them to new levels of effectiveness
- CORE data collected on over 1,000 patients shows human givens therapy to be reliably effective for the majority of patients in only a small number of sessions
- Flexible, part-time courses held in London, Bristol and York

THE HUMAN GIVENS DIPLOMA COURSE, TAUGHT BY MINDFIELDS COLLEGE, IS ALSO ACCREDITED BY NOTTINGHAM TRENT UNIVERSITY AS PART OF THE COLLABORATIVE MA PROGRAMME IN HUMAN GIVENS PSYCHOTHERAPY TAUGHT BY THE COLLEGE & NTU.

For further information about both courses, please call Kathy Hardy (College Registrar) on
01323 811440
quoting 'PYdip' or visit:
www.mindfields.org.uk

SUBSCRIPTIONS AND FEES 2010

Core Subscriptions (for various grades of membership)	Direct Debit (£)	Non-DD (£)
Members: Fellows, Associate Fellows and Graduate Members	105.00	109.00
Graduate Members during first three years after eligibility on application to the Trustees	51.00	55.00
Graduate Members on confirmation of being a <i>bona fide</i> postgraduate student with a level of income below that necessary to pay income tax	24.00	28.00
Student Subscribers	17.00	21.00
Student Subscribers earning taxable salary	51.00	55.00
Affiliates (not receiving <i>The Psychologist</i>)	20.00	24.00
Affiliates (receiving <i>The Psychologist</i>)	60.00	64.00

Division Fees	Grade	£
Division of Educational and Child Psychology	Full In training	45.00 25.00
Scottish Division of Educational Psychology	Full General In training	30.00 30.00 15.00
Division of Clinical Psychology	Full Full [first 2 years after qualification] General In training	85.00 32.00 21.00 16.00
Division of Occupational Psychology	Full In training General	40.00 20.00 15.00
Division of Forensic Psychology	Full In training General	35.00 20.00 15.00
Division of Counselling Psychology	Full Full [first 2 years after qualification] General In training	45.00 35.00 35.00 12.00
Division for Teachers and Researchers in Psychology	Full In training General	15.00 15.00 15.00
Division of Health Psychology	Full In training General	30.00 9.00 9.00
Division of Neuropsychology	Full Practitioner Full Affiliate	30.00 30.00 15.00
Division of Sport and Exercise Psychology	Full General	35.00 15.00

Section Fees	Student Rate (£)	All other (£)
Psychotherapy Section	7.50	15.00
Psychology of Education Section	2.00	10.00
Social Psychology Section	5.00	15.00
Mathematical, Statistical and Computing Section	0.00	4.00
Developmental Psychology Section	5.00	10.00
Cognitive Psychology Section	5.00	5.00
Psychobiology Section	5.00	10.00
History and Philosophy of Psychology Section	3.00	12.00
Psychology of Women Section	5.00	15.00
Transpersonal Psychology Section	3.00	10.00
Consciousness and Experiential Psychology Section	4.00	10.00
Psychology of Sexualities Section	5.00	10.00
Qualitative Methods in Psychology Section	5.00	10.00

Journal Volume (all parts for the year)	Student rate £17	All others £24
British Journal of Psychology		
Psychology and Psychotherapy: Theory, Research and Practice		
British Journal of Educational Psychology		
British Journal of Social Psychology		
British Journal of Clinical Psychology		
British Journal of Mathematical and Statistical Psychology		
Journal of Occupational and Organizational Psychology		
British Journal of Developmental Psychology		
Legal and Criminological Psychology		
British Journal of Health Psychology		
Journal of Neuropsychology		

Special Group Fees	Grade	£
Special Group for Psychologists and Social Services	Member Affiliate	15.00 5.00
Special Group in Coaching Psychology	Member Affiliate	8.50 8.50